

Propose economical and stable lepton mass matrices with texture zeros

Shu-jun Rong*

Department of Physics, Shaanxi University of Technology, Hanzhong, Shaanxi 723000, China

Abstract

There are many viable combinations of texture zeros in lepton mass matrices. We propose an economical and stable mass texture. Analytical and numerical results on mixing parameters and the effective mass of neutrinos are obtained. These results satisfy new constraints from neutrinos oscillation experiments and cosmological observations. Our proposition reveals that in the complex forest of neutrinos mixing models, a simple and robust one is still possible.

PACS numbers: 14.60.St, 14.60.Pq,

*Electronic address: rongshj@sntu.edu.cn

I. INTRODUCTION

Neutrinos oscillation is one of most mysterious phenomena in particle physics, which reveals that neutrinos are massive and mixing structure of them are nontrivial. However, how to interpret the origin of masses and special mixing pattern of neutrinos is still an open question. As useful phenomenological scenarios, special textures of lepton mass matrices and corresponding flavor groups are introduced to predict mixing parameters of neutrinos. In particular, lepton mass matrices with texture zeros [1–8] or structures alike [9–12] are widely employed in mixing models of neutrinos, which can be realized in Abelian or non-Abelian flavor groups, see Refs [13–15] for example. As an important progress, the recent Ref[16] has performed a complete survey of possible combinations of texture zeros in mass matrices of charged leptons and Majorana or Dirac neutrinos. There are so many types of possible mass matrices with texture zeros that we don't know what is the true texture chosen by nature. In order to evaluate importance of classes of texture zeros, we propose two criteria: First, the mass matrices with texture zeros should be as economical as possible; Second, predictions of texture-zeros models should be stable under perturbations of mass matrices. We consider that an economical model of texture zeros can give clear dependences of observables on parameters of lepton mass matrices. And texture-zeros models with robust predictions are more sustainable, because they can react to the progress of neutrinos experiments by introduction small modifications or perturbations.

In this letter, we propose a special combination of texture zeros of leptons mass matrices with an economical and stable structure. The mass matrix of charged leptons is Hermitian with Fritzsch texture [17–19], i.e.,

$$M_l = \begin{pmatrix} 0 & A_l & 0 \\ A_l^* & 0 & B_l \\ 0 & B_l^* & C_l \end{pmatrix}, \quad (1)$$

where C_l is real and A_l, B_l are complex parameters. And the mass matrix of Majorana neutrinos with one zero, i.e.,

$$M_\nu = \begin{pmatrix} a + c & b & 0 \\ b & a & c \\ 0 & c & a + b \end{pmatrix}, \quad (2)$$

where a is complex and b, c are real parameters. As for M_l , because of its special structure, resorting to a diagonal phase matrix, it can be transformed to a real symmetric matrix with three parameters determined by masses of charged leptons. So it can be diagonalized by a matrix with

a simple analytical expression. As for M_ν , it is magic, which predicts a trimaximal mixing pattern (TM₂) [20] if the mass matrix of charged leptons is diagonal. The structure of M_ν is characterised by the ratio b/c . So it can be diagonalized by a matrix with one real parameter. Furthermore, if a is a real parameter, M_ν would be more economical. However, in this case, ranges of mixing parameters would be smaller. So we introduce a as a complex parameter in this paper.

In following sections, the analytical expression of the lepton mixing matrix is given. Numerical predictions of the mixing parameters are obtained by scanning parameter space of mass matrices. Stabilities of these predictions are examined by introduction of random perturbations that break texture zeros of lepton mass matrices. Finally, a summary is presented.

II. ANALYTICAL FRAMEWORK

In this section, we consider the analytical expression of mixing matrix and dependence of observables on parameters in lepton mass matrices.

A. Analytical expression of mixing matrix

The lepton mixing matrix with the standard parametrization is expressed as

$$U = \text{diag}\left(e^{i\delta_e}, e^{i\delta_\mu}, e^{i\delta_\tau}\right) \cdot V \cdot \text{diag}\left(e^{-i\varphi_1/2}, e^{-i\varphi_2/2}, 1\right) \quad (3)$$

where δ_e, μ, τ are unphysical phases, $\varphi_{1,2}$ are Majorana phases, V is written as [21]

$$V = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{13}s_{23}e^{i\delta} & c_{12}c_{23} - s_{12}s_{13}s_{23}e^{i\delta} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}s_{13}c_{23}e^{i\delta} & -c_{12}s_{23} - s_{12}s_{13}c_{23}e^{i\delta} & c_{13}c_{23} \end{pmatrix}, \quad (4)$$

where $s_{ij} \equiv \sin \theta_{ij}$, $c_{ij} \equiv \cos \theta_{ij}$, δ is the Dirac CP-violating phase. This standard mixing matrix can be obtained through $U = U_l^\dagger U_\nu$, where U_l in our model comes from diagonalization of the Hermitian mass matrix M_l , i.e.

$$U_l^\dagger M_l U_l = \text{diag}\left(m_e, m_\mu, m_\tau\right), \quad (5)$$

U_ν comes from diagonalization of mass matrix of Majorana neutrinos, i.e.

$$U_\nu^T M_\nu U_\nu = \text{diag}\left(m_1, m_2, m_3\right). \quad (6)$$

In detail, $U_l = P_l O_l$, where

$$P_l = \text{diag} \left(e^{i\alpha}, e^{i\beta}, 1 \right), \quad (7)$$

where $\alpha = \arg[A_l] + \arg[B_l]$, $\beta = \arg[B_l]$, elements of O_l are written as [22, 23]

$$\begin{aligned} (O_l)_{11} &= \left[\frac{x_l - z_l}{(1 + x_l)(1 - z_l)(x_l - z_l + x_l z_l)} \right]^{1/2}, \\ (O_l)_{12} &= -i \left[\frac{(1 + z_l)x_l^3}{(1 + x_l)(x_l + z_l)(x_l - z_l + x_l z_l)} \right]^{1/2}, \\ (O_l)_{13} &= \left[\frac{(1 - x_l)z_l^3}{(1 - z_l)(x_l + z_l)(x_l - z_l + x_l z_l)} \right]^{1/2}, \\ (O_l)_{21} &= \left[\frac{(x_l - z_l)}{(1 + x_l)(1 - z_l)} \right]^{1/2}, \\ (O_l)_{22} &= i \left[\frac{x_l(1 + z_l)}{(1 + x_l)(x_l + z_l)} \right]^{1/2}, \\ (O_l)_{23} &= \left[\frac{z_l(1 - x_l)}{(1 - z_l)(x_l + z_l)} \right]^{1/2}, \\ (O_l)_{31} &= - \left[\frac{x_l z_l(1 - x_l)(1 + z_l)}{(1 + x_l)(1 - z_l)(x_l - z_l + x_l z_l)} \right]^{1/2}, \\ (O_l)_{32} &= -i \left[\frac{z_l(1 - x_l)(x_l - z_l)}{(1 + x_l)(x_l + z_l)(x_l - z_l + x_l z_l)} \right]^{1/2}, \\ (O_l)_{33} &= \left[\frac{x_l(1 + z_l)(x_l - z_l)}{(1 - z_l)(x_l + z_l)(x_l - z_l + x_l z_l)} \right]^{1/2}, \end{aligned} \quad (8)$$

where $x_l = m_e/m_\mu$, $z_l = m_e/m_\tau$. Because mass hierarchy of charged leptons is strong, approximation of O_l is

$$O_l \approx \begin{pmatrix} 1 & -ix_l^{1/2} & z_l^{3/2}x_l^{-1} \\ x_l^{1/2} & i & z_l^{1/2}x_l^{-1/2} \\ -z_l^{1/2} & -iz_l^{1/2}x_l^{-1/2} & 1 \end{pmatrix}. \quad (9)$$

Furthermore, mass parameters of M_l could also be expressed with masses of charged leptons, i. e., [22]

$$|A_l| = \left(\frac{m_e m_\mu m_\tau}{m_e - m_\mu + m_\tau} \right)^{1/2},$$

$$|B_l| = \left[\frac{(m_e - m_\mu)(m_\mu - m_\tau)(m_e + m_\tau)}{m_e - m_\mu + m_\tau} \right]^{1/2},$$

$$C_l = m_e - m_\mu + m_\tau. \quad (10)$$

As for neutrinos, $U_\nu = O_\nu P_\nu$, where

$$O_\nu = \begin{pmatrix} O_{1\nu} & O_{2\nu} & O_{3\nu} \end{pmatrix}, \quad (11)$$

$$O_{1\nu} = N_{1\nu} \begin{pmatrix} -1 + x_\nu - \sqrt{1 - x_\nu + x_\nu^2} \\ x_\nu + \sqrt{1 - x_\nu + x_\nu^2} \\ 1 \end{pmatrix}, \quad O_{2\nu} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \quad O_{3\nu} = N_{3\nu} \begin{pmatrix} -1 + x_\nu + \sqrt{1 - x_\nu + x_\nu^2} \\ x_\nu - \sqrt{1 - x_\nu + x_\nu^2} \\ 1 \end{pmatrix}, \quad (12)$$

where $x_\nu = b/c$, $N_{1\nu}$ and $N_{3\nu}$ are normalization factors expressed as

$$N_{1\nu} = \left[\sqrt{1 + (-1 + x_\nu - \sqrt{1 - x_\nu + x_\nu^2})^2 + (x_\nu + \sqrt{1 - x_\nu + x_\nu^2})^2} \right]^{-1},$$

$$N_{3\nu} = \left[\sqrt{1 + (-1 + x_\nu + \sqrt{1 - x_\nu + x_\nu^2})^2 + (x_\nu - \sqrt{1 - x_\nu + x_\nu^2})^2} \right]^{-1}. \quad (13)$$

And the diagonal phase matrix P_ν is expressed as

$$P_\nu = \text{diag} \left(e^{-i\phi_1/2}, e^{-i\phi_2/2}, e^{-i\phi_3/2} \right), \quad (14)$$

where $\phi_1 = \arg[a - \sqrt{b^2 - bc + c^2}]$, $\phi_2 = \arg[a + b + c]$, $\phi_3 = \arg[a + \sqrt{b^2 - bc + c^2}]$. Employing expressions of O_l , P_l , O_ν , and P_ν listed above, elements of the lepton mixing matrix can be written as

$$U_{ij} = (O_l^+)_{il}(O_\nu)_{1j}e^{-i(\phi_j/2+\alpha)} + (O_l^+)_{i2}(O_\nu)_{2j}e^{-i(\phi_j/2+\beta)} + (O_l^+)_{i3}(O_\nu)_{3j}e^{-i\phi_j/2}. \quad (15)$$

Obviously, ϕ_{js} are just relevant to Majorana phases of the mixing matrix.

B. Dependence of observables on parameters in lepton mass matrices

The mixing angles of leptons can be obtained from comparison of expression of U_{ij} in Eq. 15 with the form of U in the standard parametrization in Eq. 3 and Eq. 4. In detail, for θ_{13} ,

$$\sin \theta_{13} = |U_{13}| = |(O_l^+)_{11}(O_\nu)_{13}e^{-i\alpha} + (O_l^+)_{12}(O_\nu)_{23}e^{-i\beta} + (O_l^+)_{13}(O_\nu)_{33}|. \quad (16)$$

Because O_l is dependent on masses of charged leptons that are determined by experiments very well, θ_{13} in fact is dependent on three real effective mass parameter: α , β , and x_ν . The same observation can also hold for θ_{23} , θ_{13} and δ . Therefore, three mixing angles and Dirac CP violating phase are not independent to each.

As for masses of neutrinos, i.e.,

$$m_1 = |a - \sqrt{b^2 - bc + c^2}|, \quad m_2 = |a + b + c|, \quad m_3 = |a + \sqrt{b^2 - bc + c^2}|, \quad (17)$$

they are dependent on four real parameters of M_ν . And the effective mass of neutrinoless double-beta decay, i.e.,

$$m_{ee} = m_1 U_{11}^2 + m_2 U_{12}^2 + m_3 U_{13}^2, \quad (18)$$

is dependent on all 6 real parameters in M_l and M_ν .

III. PREDICTIONS OF LEPTON MASS MATRICES AND THEIR STABILITIES

A. Predictions of lepton mass matrices

We take masses of charged leptons from Particle Data Group [24]:

$$m_e = 0.51 \text{ MeV}, \quad m_\mu = 105.658 \text{ MeV}, \quad m_\tau = 1776.86 \text{ MeV}. \quad (19)$$

Then using the Mixing-Parameters-Tools package [25, 26], we scan 6 real parameters in lepton mass matrices, i.e., $\arg[A_l]$, $\arg[B_l]$, $\arg[a]$, $|a|$, b , c . The numerical results are shown in Fig. 1, including predictions for mixing angles (θ_{13} , θ_{12} , θ_{23}), Dirac CP violating phase δ , mass ratios (m_1/m_2 , m_2/m_3), the mass of the lightest neutrino m_1 , sum of masses of neutrinos $\sum m_i$, the effective mass of neutrinoless double-beta decay m_{ee} . Some comments are given as follows:

- Different from the case where both of mass matrices of charged leptons and neutrinos are of the form of Fritzsch texture [27], the octant of θ_{23} is uncertain in our case.
- Because of the strong mass hierarchy of charged leptons, the range of the Dirac CP violating phase δ is narrow, i.e., $0^\circ \pm 50^\circ$ or $180^\circ \pm 50^\circ$.
- The upper limit of the mass of the lightest neutrino is small, i.e., about 23.5 meV. So is that of the effective mass of double-beta decay i.e., about 24 meV. Accordingly, the rate of neutrinoless double-beta decay may be too small to detect in the present stage [29].
- The mass ordering of neutrinos is normal. And the hierarchy of them is not strong. so the upper limit of the sum of masses of neutrinos is also small, i.e. approximating 102 meV, which satisfies new stringent constraints from cosmological observations, e.g., $\sum m_i < 146 \text{ meV}$ at the 2σ level in Ref. [30].




FIG. 1: Predictions for mixing angles (θ_{13} , θ_{12} , θ_{23}), Dirac CP violating phase δ , mass ratios (m_1/m_2 , m_2/m_3), the mass of the lightest neutrino m_1 , sum of masses of neutrinos $\sum m_i$, the effective mass of neutrinoless double-beta decay m_{ee} . Here the unit of angles is degree.

B. Stabilities of predictions under perturbations breaking zeros

The above predictions are obtained on the basis of the special combinations of texture zeros of leptons mass matrices. In order to examine stabilities of these predictions, we introduce perturbations that break zeros in the mass matrices. For charged leptons, the perturbation is proposed

as

$$M_l = \begin{pmatrix} \epsilon_{11} & A_l & \epsilon_{13} \\ A_l^* & \epsilon_{12} & B_l \\ \epsilon_{13}^* & B_l^* & C_l \end{pmatrix}, \quad (20)$$

where ϵ_{11} and ϵ_{12} are real perturbations of order $O(0.01m_e)$, ϵ_{13} is a complex perturbation of order $O(0.01m_\mu)$. For neutrinos, the mass matrix after the perturbation is

$$M_\nu = \begin{pmatrix} a + c & b & \epsilon' \\ b & a & c \\ \epsilon' & c & a + b \end{pmatrix}, \quad (21)$$

where ϵ' is a real perturbation of order $O(0.01|a|)$. Together with initial parameters in lepton mass matrices, these parameters of perturbations are scanned. And plots of correlations of neutrinos oscillation parameters before and after perturbations are shown in Fig. 2. Some comments are given as follows:

- For θ_{12} and θ_{13} , variations brought by perturbations approximate 1° , while for θ_{23} and δ variations are nearly negligible.
- For the mass squared difference $\delta m_{21}^2 = m_2^2 - m_1^2$, the relative variation brought by perturbations approximate 10%. The relative variation of $\delta m_{31}^2 = m_3^2 - m_1^2$ is about approximate 1%.
- The mixing parameters after perturbations are still in the 3σ ranges of the experiments fit, except marginal data of θ_{13} and δm_{21}^2 . Therefore, the predictions of lepton mass matrices of our special texture are stable under random perturbations that break zeros in them.

IV. SUMMARY

There are many viable combinations of texture zeros in lepton mass matrices. According to the criteria that the structure of leptons mass matrices should be economical and the prediction of them should be robust, a special combination of texture zeros of mass matrices is proposed. The analytical expression of lepton mixing matrix is obtained, which shows clear dependence of a observable on model parameters. By scanning of parameter-space of mass matrices of leptons, numerical predictions for mixing angles, Dirac CP violating phase, and masses and effective masses of neutrinos are given. These results satisfy constraints from neutrinos oscillation experiments and those from new cosmological observations. These results are also stable under perturbations that




FIG. 2: Correlations of neutrinos mixing parameters before and after perturbations. Here the superscripts "0" in the plots denote initial mixing parameters (before perturbations). The initial mixing parameters are in the ranges of 3σ from the Ref. [28] And the unit of angles is degree.

break zeros in mass matrices. So, in the complex forest of neutrinos mixing models, a simple and robust one is still possible.

Acknowledgments

The author thank Y.F. Li for helpful discussion. This work is supported by the research foundation of Shaanxi University of Technology under the Grant No. SLGQD-13-10 and the National

- [1] P. H. Frampton, S. L. Glashow and D. Marfatia, *Zeroes of the neutrino mass matrix*, *Phys. Lett. B* **536**(2002) 79
- [2] Zhi-zhong Xing, *Texture zeros and Majorana phases of the neutrino mass matrix*, *Phys. Lett. B* **530** (2002) 159
- [3] S. Dev, S. Kumar, S. Verma and S. Gupta, *Phenomenology of two-texture zero neutrino mass matrices*, *Phys. Rev. D* **76** (2007) 013002
- [4] G. C. Branco, D. Emmanuel-Costa, R. González Felipe and H. Serôdio, *Weak basis transformations and texture zeros in the leptonic sector*, *Phys. Lett. B* **670** (2009) 340
- [5] W. Grimus and P. O. Ludl, *Maximal atmospheric neutrino mixing from texture zeros and quasi-degenerate neutrino masses*, *Phys. Lett. B* **700** (2011) 356
- [6] S. Kumar, *Implications of a class of neutrino mass matrices with texture zeros for non-zero θ_{13}* , *Phys. Rev. D* **84** (2011) 077301
- [7] D. Meloni and G. Blankenburg, *Fine-tuning and naturalness issues in the two-zero neutrino mass textures*, *Nucl. Phys. B* **867** (2013) 749
- [8] D. Meloni, A. Meroni and E. Peinado, *Two-zero Majorana textures in the light of the Planck results*, *Phys. Rev. D* **89** (2014) 053009
- [9] S. Kaneko, H. Sawanaka, M. Tanimoto, *Hybrid textures of neutrinos*, *JHEP* **0508** (2005) 073
- [10] S. Dev, S. Verma, S. Gupta, *Phenomenological Analysis of Hybrid Textures of Neutrinos*, *Phys. Lett. B* **687** (2010) 53
- [11] B. C. Chauhan, J. Pulido, M. Picariello, *Neutrino mass matrices with vanishing determinant*, *Phys. Rev. D* **73** (2006) 053003
- [12] E. I. Lashin, N. Chamoun, *Zero minors of the neutrino mass matrix*, *Phys. Rev. D* **78** (2008) 073002
- [13] W. Grimus, A. S. Joshipura, L. Lavoura, M. Tanimoto, *Symmetry realization of texture zeros*, *Eur. Phys. J. C* **36** (2004) 227-232
- [14] C. I. Low, *Abelian family symmetries and the simplest models that give $\theta_{13} = 0$ in the neutrino mixing matrix*, *Phys. Rev. D* **71** (2005) 073007
- [15] M. Hirsch, Anjan S. Joshipura, S. Kaneko, J. W. F. Valle, *Phys. Rev. Lett* **99** (2007) 151802
- [16] P. O. Ludl, W. Grimus, *A complete survey of texture zeros in the lepton mass matrices*, *JHEP* **07** (2014)

- [17] H. Fritzsch, *Weak Interaction Mixing in the Six - Quark Theory*, *Phys. Lett. B* **73** (1978) 317
- [18] H. Fritzsch, Zhi-zhong Xing, *Relating the neutrino mixing angles to a lepton mass hierarchy*, *Phys. Lett. B*, **682** (2009) 220
- [19] H. Fritzsch, *Neutrino Masses and Flavor Mixing*, *Mod. Phys. Lett. A* **30** (2015) 1530012
- [20] C. S. Lam, *Mass Independent Textures and Symmetry*, *Phys. Rev. D* **74** (2006) 113004
- [21] J. Beringer et al. (Particle Data Group), *Review of particle physics* , *Phys. Rev. D* **86** (2012) 010001
- [22] H. Fritzsch, S. Zhou, *Neutrino Mixing Angles from Texture Zeros of the Lepton Mass Matrices*, *Phys. Lett. B* **718** (2013) 1457
- [23] Zhi-zhong Xing, *Hierarchical neutrino masses and large mixing angles from the Fritzsch texture of lepton mass matrices*, *Phys. Lett. B* **550** (2002) 178
- [24] K. A. Olive et al. (Particle Data Group), *Review of Particle Physics* , *Chin. Phys. C*, **38** (2014) 090001
- [25] <http://reapmpt.hepforge.org/>
- [26] S. Antusch, J. Kersten, M. Lindner, M. Ratz, M. A. Schmidt, *Running Neutrino Mass Parameters in See-Saw Scenarios*, *JHEP* **03**(2005)024
- [27] M. Fukugita, Y. Shimizu, M. Tanimoto, T. T. Yanagida, θ_{13} in *Neutrino Mass Matrix with the Minimal Texture*, *Phys. Lett. B* **716** (2012) 294
- [28] M. C. Gonzalez-Garcia, Michele Maltoni and Thomas Schwetz, *Updated fit to three neutrino mixing: status of leptonic CP violation*, *JHEP* **11** (2014) 052
- [29] S. Dell’Oro, S. Marcocci, F. Vissani, *New expectations and uncertainties on neutrinoless double beta decay*, *Phys. Rev. D* **90** (2014) 033005
- [30] N. Palanque-Delabrouille et al., *Constraint on neutrino masses from SDSS-III/BOSS Ly α forest and other cosmological probes*, *JCAP* **02** (2015) 045